




## Production advice ware potatoes TRADITIONAL FRESH

- Big size tubers
- Uniform tuber size and shape
- High yield
- Good cooking quality
- Good resistance to common scab



### Agronomic characters

Maturity	58	Medium late
Yield early	115	Very high
Dormancy	84	Very long
Yield mature	115	Very high
Tuber size	86	Very large
Tuber shape		Oval
Number of tubers		6-8
Skin colour		Yellow
Flesh after cooking		Light yellow
Cooking type		AB - Slightly firm
Dry matter content/Starch		17,5% / 11,7%
UWW/Specific gravity		316 / 1,067
Internal bruising	12	Quite sensitive
Metribuzin sensitivity	78	●●●●●○
Little Potato disorder	52	●●○○○○



Skin and flesh colour



Cooking type



Maturity

### Plant populations

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	51.000	26	22
35/45	48.000	28	23
35/55	46.000	29	24
50/55	43.000	31	26
55/65	41.500	32	27

### Resistances

Foliage Blight	58	●●●●○○
Tuber Blight	66	●●●●○○
Alternaria	83	●●●●●●
Common scab	68	●●●●○○
Powdery scab	58	●●●●○○
Spraing	85	●●●●○○
PVY	38	●○○○○○
Yntn tuber tolerance	82	●●●●○○

#### PCN Resistance

Ro1/4	9	●●●●●●
Ro2/3	*1	●○○○○○
Pa2	*1	●○○○○○
Pa3	*1	●○○○○○

#### Wart disease

F1	10	●●●●●●
F2	*8	●●●●●●
F6		○○○○○○
F18		○○○○○○

\* HZPC own analysis/no official analysis

### Fertilizer

- Adapt fertilization to soil analysis.
- Always refer to the local and current rules about crop fertilization.
- Apply 2/3 before planting and 1/3 as top dressing.
- Nitrogen (N) : Very high inputs (>250Kg N/Ha inclusive of soil supply)
- Potassium and phosphate fertilization as standard advice.
- To be able to harvest early, organic manure is not advised.



## Production advice ware potatoes **TRADITIONAL FRESH**



### Pre-treatment and planting

- FABULA has a long dormancy.
- The apical dominance is quite strong.
- A heat treatment is preferred to break the dormancy.
- De-sprouting will increase the risk of little potato disorder.
- Sensitive to little potato disorder.
- The best results are obtained on lighter soils.
- Plant into warm soils, don't plant too early.
- Do not plant in cold soils as this will increase the risk of irregular emergence.




### Growing attention points

- Emergence is somewhat slow, later on foliage develops well.
- Tuber number is quite low and tubers easily get too big.
- Preventive spraying against Phytophthora is advised.



### Haulm killing and harvest

- The skin should be fully set before harvest.
- FABULA is moderately susceptible to bruising. Reduce drop heights and avoid mechanical damage.
- Prevent mechanical damage to improve storability.



### Storage

- Prevent dehydration, cool with a small difference between cooling air and product temperature.
- Take care of the wound healing period.
- Dry quickly after harvest and keep dry to help prevent silver scurf.
- Storage temperature: 5 °C.
- Stable temperature during storage prevents silver scurf development.